

Membrane Structural Alterations in Murine Stratum Corneum: Relationship to the Localization of Polar Lipids and Phospholipases

To the Editor:

The principal theme of the article by Elias et al [1] is that the stratum corneum is not a metabolically inert tissue. However, this is not a new concept, and the report [1] merely adds an additional enzyme to the existing list of activities found in stratum corneum [2-4].

In support of their major point, the authors demonstrate that very small amounts of phospholipids can be detected in inner but not outer stratum corneum, as previously reported by others [5,6]. The present workers removed the outer stratum corneum material by tape stripping, which raises a question as to whether the tape adhesives interfered with the lipid analyses.

They also demonstrate, in Fig 1a of their paper, that the inner stratum corneum stains for phospholipids while the outer stratum corneum does not. This same figure was published nine years earlier (Fig 7 of Ref 7) and said to demonstrate that only the *viable* epidermis stained for phospholipids, while the stratum corneum did not stain [7]. What new information led to the present reinterpretation?

The authors demonstrate phospholipase activity in the intercellular spaces throughout the stratum corneum, and suggest that "First, increased generation of fatty acids from phospholipids, because of their amphipathic characteristics, may facilitate the dispersion of lamellar lipids into micelles within the outermost layers of the stratum corneum, hence, leading to desquamation." This proposal would seem to be irreconcilable with the data in Table II of their paper, which indicate that phospholipids could not be detected in the outer stratum corneum, and with the statement that the stacks of membrane bilayers in the outer stratum corneum can be revealed by osmium vapor fixation. Two references [8,9] were cited to support the latter point, but examination of those two references reveals that the lamellae were not localized to any particular region of the stratum corneum. In three other recent papers [10-12], the senior author of the present article [1] stated that the lamellae disappeared or could not be found beyond the mid-to-outer-third of the stratum corneum, which is contradictory to the present statement.

Finally, three recent publications in this journal [13-15] have demonstrated that postfixation with RuO_4 permits the visualization of lipid lamellae throughout the stratum corneum and even in desquamated material. It is curious that the authors of the article [1] should fail to cite these relevant publications.

Philip W. Wertz, Ph.D.
Donald T. Downing, Ph.D.
Donald C. Swartzendruber, Ph.D.
Kathi C. Madison, M.D.
John S. Strauss, M.D.
Marshall Dermatology Research Laboratories
University of Iowa College of Medicine
Iowa City, Iowa

REFERENCES

- Elias PM, Menon GK, Grayson S, Brown BE: Membrane structural alterations in murine stratum corneum: Relationship to the localization of polar lipids and phospholipases. *J Invest Dermatol* 91:3-10, 1988
- Barrett JG, Scott IR: Pyrrolidone carboxylic acid synthesis in guinea pig epidermis. *J Invest Dermatol* 81:122-124, 1983

- De Young L, Ballaron S, Epstein W: Transglutaminase activity in human and rabbit ear comedogenesis: a histochemical study. *J Invest Dermatol* 82:275-279, 1984
- Baden HP, Hooker PA, Kubilus J, Tarascio A: Sulfatase activity of keratinizing tissues in X-linked ichthyosis. *Ped Res* 14:1347-1348, 1980
- Gray GM, White RJ, Williams RH, Yardley HJ: Lipid composition of the superficial stratum corneum of pig epidermis. *Br J Dermatol* 106:59-63, 1982
- Bowser PA, White RJ: Isolation, barrier properties and lipid analyses of stratum compactum, a discreet region of the stratum corneum. *Br J Dermatol* 112:1-14, 1985
- Elias PM, Brown BE, Fritch P, Goerke J, Gray GM, White RJ: Localization and composition of lipids in neonatal mouse stratum granulosum and stratum corneum. *J Invest Dermatol* 73:339-348, 1979
- Elias PM, McNutt NS, Friend D: Membrane alterations during cornification of mammalian squamous epithelia: A freeze fracture, tracer and thin section study. *Anat Rec* 189:577-593, 1977
- Elias PM, Friend D: The permeability barrier in mammalian epidermis. *J Cell Biol* 65:180-191, 1975
- Menon GK, Grayson S, Brown BE, Elias PM: Lipokeratinocytes of the epidermis of a cetacean (*Phocena phocena*): Histochemistry, ultrastructure and lipid composition. *Cell Tissue Res* 244:385-394, 1986
- Elias PM, Menon GK, Grayson S, Brown BE, Rehfeld J: Avian sebo-keratinocytes and marine mammal lipokeratinocytes: Structural, lipid biochemical, and functional considerations. *Am J Anat* 180:161-177, 1987
- Williams ML, Elias PM: The extracellular matrix of the stratum corneum: Role of lipids in normal and pathological function. *CRC Crit Rev Therapeutic Drug Carrier Systems* 3:95-122, 1987
- Madison KC, Swartzendruber DC, Wertz PW, Downing DT: Presence of intact intercellular lipid lamellae in the upper layers of the stratum corneum. *J Invest Dermatol* 88:714-718, 1987
- Swartzendruber DC, Wertz PW, Madison KC, Downing DT: Evidence that the corneocyte has a chemically bound lipid envelope. *J Invest Dermatol* 88:709-713, 1987
- Wertz PW, Swartzendruber DC, Madison KC, Downing DT: Composition and morphology of epidermal cyst lipids. *J Invest Dermatol* 89:419-425, 1987

REPLY

In response to the comments of Wertz et al about our recent paper (1), first, we agree that the observation that the stratum corneum may be metabolically active is not new; we [2-4] and others (e.g., Refs 5-7) have demonstrated various types of hydrolytic enzyme activity both biochemically and cytochemically. The relative merits of adding an additional enzyme (phospholipase in the case) ought to be judged by the potential biological significance of the enzyme in the tissue concerned. In this report we not only show the localization of the enzyme to stratum corneum (SC) interstices, but also the likely consequences; i.e., a precipitous drop in phospholipid content with resultant changes in the structure of intercellular lamellae. Thus, the importance of our observations lies in the co-localization of this enzyme activity with lipid precursors, a finding that may have implications for both the formation of a hydrophobic barrier